



Product Guide



A WORLD TO
CARE AND PROTECT

MEDICATED PREMIXES

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Ampicilina 100	Ampicillin tri-hydrate 100 mg/g	Oral administration via feed. Poultry: 2 to 5 kg/ton of feed. Calves: 5 to 10 kg/ton of feed.	Poultry and calves	1 kg, 5 kg and 25 kg	Meat: 8 days Eggs: 10 days
Coliscid	Colistin sulfate 720 000 UI/g	Oral administration via feed. 2 to 4 kg/ton of feed.	Poultry	1 kg, 5 kg and 25 kg	Meat: 7 days Eggs: Zero days
Oxitetraciclina 200	Oxytetracycline 200mg/g	Oral administration via feed. Poultry: 2,5 kg/ ton of feed. Rabbits: 2,5 kg/ton of feed.	Poultry and rabbits	1 kg, 5 kg and 25 kg	Meat and offal: 7 days Do not use in animals producers of eggs for human consumption
Tilfur Medicated Premix	Tylosin phosphate 100 mg/g	Oral Administration via feed. 1 to 2 kg/ ton of feed.	Poultry and other species	1 kg, 5 kg and 25 kg	Meat: 0 days
Tiamulol	Tiamulin hydrogen fumarate 100 mg/g	Oral administration via feed. Poultry: 3 to 4 kg/ton of feed.	Poultry and other species	1 kg, 5 kg and 25 kg	Meat: 5 days
Trimecoli	Trimethoprim 20 mg/g Sulfadimethoxine 100 mg/g	Oral administration via feed. Poultry and rabbits: 3 to 5 kg/ton of feed.	Poultry and rabbits	1 kg, 5 kg and 25 kg	Meat and offal: 6 days Eggs: 10 days
Zoomisol	Levamisole HCL 50 mg/g	Oral administration via feed. Poultry: 2 kg/ton of feed. Sheep and goats: 2 kg/ton of feed. Bovine: 2kg/ton of feed.	Poultry, cattle, sheep and goats	1 kg, 5 kg and 25 kg	Meat: 3 days Eggs: Do not use in animals producers of eggs for human consumption

WATER SOLUBLE POWDERS

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Amoxicillin 800	Amoxicillin tri-hydrate 800 mg/g	Oral administration via drinking water. Broilers and hens: The recommended dose is 8-16mg amoxicillin tri-hydrate/kg bw/day (equivalent to 10-20 mg of the medicine/kg body weight per day). Other species please contact.	Poultry	100g and 1 kg	Poultry: 1 Day Eggs: Do not use in animals producing eggs for human consumption
Coliscid WS	Colistin Sulfate 1060 000 UI/g	Oral administration via drinking water. Poultry: 75 000 U.I./kg bw/day.	Poultry	100 g and 1 kg	Meat: 7 days Eggs: Zero days
Doxycycline 500	Doxycycline Hyclate 500mg/g	Oral administration. Poultry: 25 mg doxycycline hyclate/kg bw/day (corresponding to 50 mg of product per kg bw), for 3-5 consecutive days.	Poultry	100 g and 1 kg	Meat: 5 days Eggs: Do not use in animals producing eggs for human consumption
Micoresp	Tylosin Tartrate 100mg/g Doxycycline Hyclate 100mg/g Bromhexine HCl 3,5 mg/g	Oral administration via drinking water. 2 to 3 g/L of drinking water.	Poultry	100 g and 1 kg	Meat and offal: Poultry: 7 days
Oxytetracycline 500	Oxytetracycline HCl 500 mg/g	Oral administration. Poultry: 200 mg of the medicine/L of drinking water.	Poultry	100 g and 1 kg	Meat: 7 days Eggs: do not use in animals producers of eggs for human consumption
Tilfur 1000	Tylosin Tartrate 1000mg/g	Oral administration via drinking water. Chicken and turkey: CRD - 0,5 g/L of drinking water. Necrotic Enteritis: 0,1 g/L of drinking water.	Poultry and turkey	100 g and 1 kg	Meat and offal: Poultry: 0 days Turkeys: 1 day

WATER SOLUBLE POWDERS (CONT.)

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Tricobacter	Dimetridazole 100 mg/g Sulfadimethoxine sodium 100mg/g Vitamin K3 (MSB) 5mg/g	Oral administration via drinking water. Poultry: 2 to 2,5 g/L of drinking water. Rabbits: 1,5 to 2 g/L of drinking water.	Poultry and rabbits	100 g and 1 kg	Meat: 15 days Do not use in animals producers of eggs for human consumption
Tiamulol WS	Tiamulin Hydrogen Fumarate 100mg/g	Oral administration. Poultry: 24,7 mg tiamulin hydrogenfumarate/kg bw WARNING: DO NOT ADMINISTER WITH IONOPHORES (LIKE MONENZIN, SALINOMYCIN, NARASIN)	Poultry	100 g and 1 kg	Meat: 5 days

ORAL SOLUTIONS

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Broncolis	Bromhexine HCl 5 mg/ml Eucalyptol 40 mg/ml	Oral administration via drinking water. Poultry: 0,5 to 1 ml/L of drinking water. Turkeys: 1 to 1,5 ml/L of drinking water. Sheep: 2 ml/L of drinking water. Goats: 1 ml/L of drinking water. Lambs: 1,5 ml/L of drinking water. Racing pigeons and singing birds: 1 to 3 ml/L of drinking water.	Poultry, cattle, camel, sheep, goats, racing pigeons, songbirds and ornamental birds	1 L, 5 L and 25 L	Poultry: 2 dias Cattle: 3 days Do not use in horses whose meat is intended for human consumption. Do not administer to animals whose milk is intended for human consumption.
Enroxina	Enrofloxacin 100 mg/ml	Poultry and rabbits: 10 mg of enrofloxacin/kg bw	Poultry and rabbits	1 L, 5 L and 25 L	Poultry - Meat and offal: 3 days. Turkeys - Meat and offal: 3 days Rabbits - Meat and offal: 15 days Other species - Meat and offal: 5 days

VITAMIN COMPLEXES

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Vitapro® AD3EC + K	Vitamin A, D3, E, C and K3	Oral administration via drinking water. Poultry, horses and rabbits: 0,5 ml/L of drinking water. Ruminants: 1 ml/L of drinking water.	Poultry, cattle, calves, camel, horses, sheep, goats and rabbits	1 L, 5 L and 25 L	N.A.
Vitapro® B + K	Vitamin B1, B2, B3, B6, B12, K3 Calcium Panthotenate	Oral administration via drinking water. Horses: 10 to 20 ml/animal/day/a period of 3 days. Other species: 1 to 2 ml/L of drinking water.	Poultry, cattle, calves, camel, horses, sheep, goats and rabbits	1 L, 5 L and 25 L	N.A.
Vitapro® E + Selenium	Vitamin E Sodium Selenite	Oral Administration via drinking water. For the deficiencies in vitamin E and for detoxification: Poultry: 50 ml/100 L of water during 5 days. To improve the properties of meat preservation: Poultry: 100 ml/100 L of water during 15 days before slaughtering. Turkeys: 100 ml/100 L of water during 30 days.	Poultry, cattle, camel and sheep	1 L, 5 L and 25 L	N.A.
Vitapro® Hepagen	Vitamin E, Vitamin B1, B2, B3, B6, B12 Calcium Panthotenate Inositol Choline Chloride Methionine	Oral administration via drinking water. Calves: 2 to 8 ml/100 kg b.w. Other species: 1 to 2 ml/L of drinking water.	Poultry, cattle, calves, camel, horses, sheep, goats and rabbits	1 L, 5 L and 25 L	N.A.
Vitapro® Multi	Vitamin A, D3, E, B1, B2, B3, B6, B12, K3, H, Calcium Panthotenate Choline Chloride L-treonine L-tryptofane	Oral administration via drinking water. Poultry: 2 to 3 ml/L of drinking water. Sheep and goats: 5 ml/10 kg body weight. Rabbits: 1 to 2 ml/L of drinking water. Cattle, calves and horses: 10 to 20 ml/100 kg body weight.	Poultry, cattle, calves, camel, horses, sheep, goats and rabbits.	1 L, 5 L and 25 L	N.A.
Vitapro® Amino L	Vitamin A, D3, E, B1, B2, B3, B6, B12, K, H L-carnitine Calcium Panthotenate Choline Chloride Lysine Methionine Aminoacid Pool	Oral administration via drinking water. Horses: 5 to 20 ml/100 kg b.w. Other species: 1 to 2 ml/L of drinking water.	Poultry, cattle, calves, camel, horses, sheep, goats and rabbits	1 L, 5 L and 25 L	N.A.

VITAMIN AND MINERAL CORRECTORS

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Foscavit Ferti Plus	Vitamin A, D3, E, B1, B3 Iron Iodine Cobalt Copper Manganese Zinc (oxide + chelate) Selenium Saccharomyces Cerevisiae	200 g/cow/day - in the 3 first months of lactation Until 25 L/day - 150 g/cow/day 25 - 35 L/day - 180 g/cow/day More than 35 L/day - 200 g/cow/day	Dairy cows	25 kg	N.A.

VITAMIN AND MINERAL CORRECTORS (CONT.)

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Foscavit® T Powder	Vitamins A,D3, E, B1, B2, B3, B6,B12 Iron Manganese Copper Selenium Zinc Iodine Cobalt	10 g/kg of feed.	Poultry, cattle, calves, camel, horses, foals, sheep, goats and rabbits	1 kg, 5 kg and 25 kg	N.A.
Foscavit® T Oral Suspension	Vitamins A, D3,E, B1, B2, B3, B6, B12 Calcium Panthotenate Iron Iodine Cobalt Copper Manganese Zinc Selenium	Poultry: 2,5 ml/L of drinking water. Dairy cows: 40 to 50 ml/day or 5 ml/L of drinking water. Calves: 10 to 15 ml/day or 5 ml/L of drinking water. Steers: 7,5 to 10 ml/100 kg of body weight or 5 ml/L of drinking water. Young sheep and goats: 2,5 ml/L of drinking water.	Poultry, cattle, calves, camel, sheep and goats	1 L and 5 L	N.A.

DIETETIC FEED

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Avisstress® Water Soluble Powder	Vitamins A, D3, E, B1, B2, B3, B6, B12, K3 L-carnitine Calcium Pantothenate	2,5 to 5 g/L of drinking water.	Poultry, cattle, calves, camel, horses, sheep, goats, rabbits and other species	100 g and 1 kg	N.A.
Cetosil® SG Oral Syrup	Choline chloride Niacin Cobalt Propylene glycol Glycerol Sorbitol Magnesium	Dairy cows : 5,5 dl/cow/day. Sheep and goats : 30 to 50 ml/cow/day.	Dairy cows, sheep and goats	1 L and 5 L	N.A.
Glical Oral Syrup	Calcium Cobalt Copper Manganese Zinc Glycerol Propylene Glycol Sorbitol	500 ml after the firsts signs of parturition. 500 ml right after parturition. 500 ml 12-24h after parturition.	Dairy cows	500 ml	N.A.
Mag-bicar	Sodium Magnesium Calcium	Dairy cows: 150 to 200 g/cow/day. Calves: 120 to 160 g/cow/day. Sheep and goats: 20 to 40 g/animal/day.	Dairy cows, sheep and goats	25 kg	N.A.

COMPLEMENTARY FEED

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Bio-Sac® Powder	<i>Bacillus subtilis</i> (ATCCPTA-6737) Calcium Formate	10 g/kg of feed	Poultry, cattle, sheep, camel, goats, horses and rabbits	100 g 1 kg and 5 kg	N.A.
Bio-Sac® WS Water Soluble Powder	<i>Bacillus subtilis</i> (ATCCPTA-6737)	2 to 4 g/L of drinking water.	Poultry, cattle, sheep, camel, goats, horses and rabbits	100 g 1 kg and 5 kg	N.A.
Imuno-Rumi®	Essential oils / Functional extracts (spices) Natural selected plants Mycotoxin binders and extracts of the plant <i>Cynara Cardunculos</i> .	Dairy cows: Pre-calving: 40 g/cow/day during the 21 days. After delivery: 100 g/cow/day, up to 21 days. Lactating animal: 60 g/cow/day. Sheep and Goats: Pre-calving: 10 g/animal/day during the 21 days. After delivery: 25 g/animal/day, up to 21 days. During lactation: 25 g/animal/day. Calves and steers: Initiation: 10 g/animal of 50-100 kg/day. Growth up to 6 months: 25 g/animal/day. From 6 to 24 months: 40 g/animal/day. Mix in the feed.	Dairy cows, sheep, goats, camel and calves	25 kg	N.A.
Liporenal OS	Sodium acetate Amonium Chloride Magnesium Sulfate Methionine Sorbitol Dextrose	1 L per 1000 L of drinking water during 5 days	Poultry	1 L and 5 L	N.A.
Rehidral® Oral Solution	Potassium Chloride Sodium Glucose Citric Acid	Poultry and other birds: 3 to 6 ml/L of drinking water during 5 to 10 days and whenever necessary. Calves (diarrhea or adaptation): 50 ml/L of warm water used to prepare the milk replacer during 2 to 3 days, and more 3 days in the milk replacer. Foals: 60 ml/L of drinking water during diarrheic disorders or in periods of intense heat. Horses: 50 ml/L of drinking water as an adjuvant of the treatment for dehydration.	Poultry and other birds, calves, camels, foals and horses	1 L and 5 L	N.A.

NATURAL PRODUCTS

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Optiprobios® Carvactyme	Essential oils of <i>Origanum Vulgaris</i> <i>Tymus Vulgaris</i>	Administer 1 ml/L of drinking water during the most critical periods (first weeks of life, stressful situations, prolonged transport, trade show presence, sudden changes in temperature, change of location, etc.). Intestinal problems: 2 to 3 ml/L of drinking water.	All kind of species	1 L and 5 L	N.A.
Optiprobios® Coccid	Extract of <i>Cinnamomum Zeylanicum</i> Essencial oil of <i>Oreganum Vulagare</i> and <i>Thymus Vulgaris</i>	Administer 2 to 3 ml/L of drinking water during 5 consecutive days.	All kind of species	1 L and 5 L	N.A.
Optiprobios® Muda	Methionine Lysine Biotin Vitamins B1, B6, C, E Zinc Copper <i>Cynara Scolimus</i> extracts	Administer 5 g/L of drinking water or 7,5 g/kg of feed. Initiate the administration at least one week before the period of moulting.	Poultry and all kind of birds	1 kg, 5 kg and 25 kg	N.A.

NATURAL PRODUCTS (CONT.)

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Optiprobios® Parasitas	Carica Papaya Annona Squamosa Andrographis Paniculata	Poultry: 1,5 ml to 3 ml/L of drinking water during 3 consecutive days. Other species: 2 ml to 4 ml/L of drinking water during 3 consecutive days.	All kind of production/farm/animals	1 L and 5 L	N.A.
Optiprobios® Resp	Extracts of Glycyrrhiza Glabra Lonicera Japonica Cynara Scolimus	Administer 1 to 2 ml/L of drinking water, during 3 to 5 days. In large birds (2 to 3 kg body weight): administer 0,5 ml once or twice a day, with a pipette or syringe directly into the nozzle, during 3 to 5 days.	Poultry	1 L and 5 L	N.A.

MICOTOXIN ADSORBENTS

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Adsortox Promo	Pure extracts of Mannanoligo saccharides and Beta-Glucans provenient of soluble yeast cell walls Cynarine, a chemical that is biologically active of Cynara Scolimus Sorbitol Choline Chloride	Poultry: 2 to 3 ml/L of drinking water. Other species: 2 ml/kg body weight or 2 to 3 ml/L of drinking water.	All kind of production/farm animals	1 L and 5 L	N.A.
Adsortox Polival	Diatoms Montmorillonite/ Activated Bentonite Yeast cell walls	Mix in the feed at a rate of 2 to 5 g/kg of dry matter (of the total feed).	All kind of production/farm animals	20 kg	N.A.
Adsortox Polival Plus	Diatoms Montmorillonite/ Activated Bentonite HSCAS Yeast cell walls Cynara Scolymus extract	Mix in the feed at a rate of 0,5 to 2 kg/ton of dry matter (of the total feed).	All kind of production/farm animals	20 kg	N.A.

DISINFECTANTS

Product Name	Composition	Dosage and Administration	Target Species	Packaging	Withdrawal Period
Clorhexidina 1%	Chlorhexidine diglucuronate	Hygiene and treatment of wounds and abscesses, topical treatment of gingivitis: chlorhexidine 1% Prevention of infections of the umbilical cord: chlorhexidine 0,5% Hygiene and treatment of ear infections: chlorhexidine 0,2% Topical treatment of candidiasis and upper respiratory tract infections: 5-15 ml of the product/ 1L of drinking water	All kind of production/farm animals	1 L and 5 L	N.A.
Zoopharmacid	Disinfectant based on Quaternary Ammonium and Aldehyde	First clean the surfaces thoroughly. Disinfect by spraying: 1% Zoopharmacid. Disinfect by fogging: 1,5 L + 3 L water Zoopharmacid/1000m ³ .	All kind of production/farm animals	1 L, 5 L and 10 L	N.A.
Zoiod	Iodine	Its concentration of application generally varies between 0,2% and 0,5%, containing approximately 110 ppm of active iodine in solution at 1%.	All kind of production/farm animals	1 L, 5 L and 10 L	N.A.



Rua da Liberdade, 77
2050-023 Aveiras de Baixo
(Azambuja) - PORTUGAL

T. (+351) 263 470 160/162
F. (+351) 263 470 169
E. export.sales@zoopharma-international.pt

www.zoopharma-international.pt

A WORLD TO
CARE AND PROTECT